

LEGENDRE TRANSFORMATION FOR REGULARIZABLE LAGRANGIANS IN FIELD THEORY

OLGA KRUPKOVÁ AND DANA SMETANOVÁ

Mathematical Institute of the Silesian University in Opava

Bezručovo nám. 13, 746 01 Opava, Czech Republic

e-mail: Olga.Krupkova@math.slu.cz, Dana.Smetanova@math.slu.cz

ABSTRACT. Hamilton equations based not only upon the Poincaré–Cartan equivalent of a first-order Lagrangian, but rather upon its Lepagean equivalent are investigated. Lagrangians which are singular within the Hamilton–De Donder theory, but regularizable in this generalized sense are studied. Legendre transformation for regularizable Lagrangians is proposed, and Hamilton equations, equivalent with the Euler–Lagrange equations, are found. It is shown that all Lagrangians affine or quadratic in the first derivatives of the field variables are regularizable. The Dirac field and the electromagnetic field are discussed in detail.

KEYWORDS. Lagrangian, Poincaré–Cartan form, Lepagean form, Hamilton–De Donder equations, Hamilton p_2 -equations, regularity, regularizable Lagrangian, Legendre transformation, Dirac field, electromagnetic field

MS CLASSIFICATION. 70G50, 58Z05

PACS NUMBERS. 02.40, 11.10

1. INTRODUCTION

If λ is a Lagrangian defined on J^1Y (the first jet prolongation of a fibred manifold $\pi : Y \rightarrow X$), and θ_λ is its Poincaré–Cartan form then the *Euler–Lagrange equations* are equations for local sections $\gamma : X \rightarrow Y$ of π as follows: $J^1\gamma^*i_\xi d\theta_\lambda = 0$, for every vertical vector field ξ on J^1Y . A geometric setting for the Hamilton theory on fibred manifolds goes back to Goldschmidt and Sternberg, who in their famous paper [7] proposed to consider *Hamilton equations* as an extension of the Euler–Lagrange equations to local

Research supported by Grants MSM:J10/98:192400002, VS 96003 and FRVŠ 1467/2000 of the Czech Ministry of Education, Youth and Sports, and Grant 201/00/0724 of the Czech Grant Agency

sections $\delta : X \rightarrow J^1Y$, namely, $\delta^*i_\xi d\theta_\lambda = 0$. To become *equivalent* with the Euler–Lagrange equations, the Lagrangian has to satisfy the *regularity condition*

$$(1.1) \quad \det\left(\frac{\partial^2 L}{\partial y_i^\sigma \partial y_k^\nu}\right) \neq 0.$$

Goldschmidt–Sternberg’s approach, now adopted as standard (cf. [2], [5], [6], [8], [15–17] and many others), however seems not to be quite satisfactory. This turns out namely if generalizations to higher order are considered, or if concrete physical fields are studied: unfortunately, almost all of them are degenerate in the sense of the regularity condition (1.1). Paralelly, Dedecker in 1977, and Krupka in 1983 considered another, from the mathematical point of view a more natural, extension of the Euler–Lagrange equations, based not upon a Poincaré–Cartan form as above, but rather upon a general *Lepagean equivalent of a Lagrangian* [3], [11]. As noticed already by Dedecker in [3], this approach opens a possibility to study “regularizations” of singular Lagrangians.

This paper develops the idea to understand Hamilton equations in the above mentioned generalized sense. However, we differ from Dedecker and Krupka in some points. First of all, we consider Lepagean equivalents which are *at most 2-contact*, i.e. of the form $\theta_\lambda +$ some auxiliary 2-contact term. Such Hamilton equations, first considered in [14] and called there *Hamilton p_2 -equations* can be viewed as a “first correction” to the standard Hamilton equations. In [14], we studied relations with the Euler–Lagrange equations, and obtained appropriate *regularity conditions*, generalizing (1.1). The aim of this paper is to propose *Legendre transformation for Hamilton p_2 -equations*, and to apply the results to concrete physically interesting first order Lagrangians, namely, Lagrangians *affine* or *quadratic* in the first derivatives of the field variables. Comparing our approach with Dedecker [3], one can see that our concept of regularity is stronger, and Legendre transformation is understood in a completely different way.

Contrary to the standard approach, where all affine and many quadratic Lagrangians are singular, we show that all these Lagrangians are regularizable, admit Legendre transformation, and provide Hamilton equations which are *equivalent* with the Euler–Lagrange equations (i.e., do *not* contain constraints). We also show that under certain additional conditions Hamilton p_2 -equations of a Lagrangian coincide with the *usual* Hamilton equations of an appropriate *equivalent Lagrangian*. We study in detail the case of the Dirac field and the electromagnetic field, and find the corresponding “corrected” momenta and Hamiltonian which could be alternatively used for (unconstrained) quantization.

Finally, we note that results and techniques presented in this paper can be generalized to higher-order variational problems [13], [18].

2. PRELIMINARIES

Throughout the paper all manifolds and mappings are smooth, and summation convention is used. We consider a fibred manifold $\pi : Y \rightarrow X$, $\dim X = n$, $\dim Y = m + n$, and its first (respectively, second) jet prolongation $\pi_1 : J^1Y \rightarrow X$ (resp. $\pi_2 : J^2Y \rightarrow X$). Natural fibred projections $J^kY \rightarrow J^lY$, where $0 \leq l < k \leq 2$, are denoted by $\pi_{k,l}$. A fibred chart on Y (respectively, associated chart on J^1Y) is denoted by (V, ψ) , $\psi = (x^i, y^\sigma)$ (respectively, (V_1, ψ_1) , where $V_1 = \pi_{1,0}^{-1}(V)$ and $\psi_1 = (x^i, y^\sigma, y_i^\sigma)$). We use the following

notations:

$$(2.1) \quad \omega_0 = dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n, \quad \omega_i = i_{\partial/\partial x^i} \omega_0, \quad \omega_{ij} = i_{\partial/\partial x^j} \omega_i, \quad \cdots,$$

and

$$(2.2) \quad \omega^\sigma = dy^\sigma - y_j^\sigma dx^j.$$

It is important to note that $(dx^i, \omega^\sigma, dy_j^\sigma)$ is a *basis of 1-forms on J^1Y* . A mapping $\gamma : X \rightarrow Y$ defined on an open subset $U \subset X$ is called a *section* of the fibred manifold π if the composite mapping $\pi \circ \gamma$ is the identity mapping of U . Quite analogously, a section of the fibred manifold π_1 is defined. Notice that a section of π_1 need not be of the form of prolongation of a section of π . Accordingly, a section δ of the fibred manifold π_1 is called *holonomic* if $\delta = J^1\gamma$ for a section γ of π .

Recall that every q -form η on J^1Y admits a unique (canonical) decomposition into a sum of q -forms on J^2Y as follows:

$$(2.3) \quad \pi_{2,1}^* \eta = h(\eta) + \sum_{k=1}^q p_k(\eta),$$

where $\pi_{2,1}$ is the canonical projection $J^2Y \rightarrow J^1Y$, $h(\eta)$ is a horizontal form, called the horizontal part of η , and $p_k(\eta)$, $1 \leq k \leq q$, is a k -contact form, called the k -contact part of η (see e.g. [9]). For our purposes it is sufficient to recall that in fibred coordinates a horizontal form on J^1Y is expressed by means of wedge products of the differentials dx^i only, with the components dependent upon the coordinates $(x^i, y^\sigma, y_j^\sigma)$. Similarly, a 1-contact (respectively, 2-contact) form contains only wedge products of the differentials dx^i with *one* (respectively, *two*) of the contact forms (2.2).

By a *first-order Lagrangian* we shall mean a horizontal n -form λ on J^1Y . This means that in every fibred chart,

$$(2.4) \quad \lambda = L\omega_0$$

where $L = L(x^i, y^\sigma, y_j^\sigma)$. A form ρ is called a *Lepagean equivalent* of a Lagrangian λ if (up to a projection) $h(\rho) = \lambda$, and $p_1(d\rho)$ is a $\pi_{2,0}$ -horizontal form [9]. All (first order) Lepagean equivalents of a Lagrangian of order one take the form

$$(2.5) \quad \rho = \theta_\lambda + \nu,$$

where θ_λ is the *Poincaré–Cartan equivalent* of λ , i.e.

$$(2.6) \quad \theta_\lambda = L\omega_0 + \frac{\partial L}{\partial y_j^\sigma} \omega^\sigma \wedge \omega_j,$$

and ν is an *arbitrary at least 2-contact n -form*, i.e. such that $h(\nu) = p_1(\nu) = 0$.

If ρ is a Lepagean equivalent of λ then the $(n+1)$ -form $E_\lambda = p_1(d\rho)$ is called the *Euler–Lagrange form* of the Lagrangian λ . Two Lagrangians λ_1 and λ_2 are called

equivalent if $E_{\lambda_1} = E_{\lambda_2}$. Recall that Lagrangians λ_1 and λ_2 are equivalent on an open set $U \subset J^1Y$ if and only if there exists an $(n-1)$ -form φ such that $\lambda_2 = \lambda_1 + h(d\varphi)$ [9].

Besides the Poincaré–Cartan equivalent (2.6), the family (2.5) of Lepagean equivalents of a Lagrangian contains another distinguished Lepagean equivalent uniquely determined by the Lagrangian, namely,

$$(2.7) \quad \rho_\lambda = L\omega_0 + \sum_{k=1}^n \left(\frac{1}{k!}\right)^2 \frac{\partial^k L}{\partial y_{j_1}^{\sigma_1} \dots \partial y_{j_k}^{\sigma_k}} \omega^{\sigma_1} \wedge \dots \wedge \omega^{\sigma_k} \wedge \omega_{j_1 \dots j_k}$$

[10] (cf. also [1]). It is called the *Krupka equivalent* of λ , and has the following important property: $d\rho_\lambda = 0$ if and only if $E_\lambda = 0$; the latter condition, however, means that $\lambda = h(d\varphi)$ (a so called trivial Lagrangian).

With the help of Lepagean equivalents of a Lagrangian one obtains an intrinsic formulation of the *Euler–Lagrange* and *Hamilton equations* as follows (cf. [9], [11]). A section γ of the fibred manifold π is an *extremal* of λ if and only if

$$(2.8) \quad J^1\gamma^* i_{J^1\xi} d\rho = 0$$

for every π -vertical vector field ξ on Y . A section δ of the fibred manifold π_1 is called a *Hamilton extremal* of ρ if

$$(2.9) \quad \delta^* i_\xi d\rho = 0,$$

for every π_1 -vertical vector field ξ on J^1Y . The equations (2.8) and (2.9) are called the *Euler–Lagrange* and the *Hamilton equations*, respectively.

Notice that while the Euler–Lagrange equations (2.8) are uniquely determined by the Lagrangian, Hamilton equations (2.9) depend upon the choice of ν . Consequently, one has many different “Hamilton theories” associated to a given variational problem.

Clearly, if γ is an extremal then $J^1\gamma$ is a Hamilton extremal; conversely, however, a Hamilton extremal need not be holonomic, and thus a jet prolongation of some extremal. This suggests a definition of regularity as follows: A Lepagean form is called *regular* if every its Hamilton extremal is holonomic [12].

Hamilton equations (2.9) where $\rho = \theta_\lambda$ (respectively, ρ is *at most 2-contact*) are called *Hamilton–De Donder equations* [4], [7] (respectively, *Hamilton p_2 -equations* [14]).

3. HAMILTON p_2 -EQUATIONS AND LEGENDRE TRANSFORMATION FOR FIRST-ORDER LAGRANGIANS

In the sequel we consider Lepagean forms (2.5) where ν is *2-contact*. Moreover, we suppose $\nu = p_2(\beta)$, where β is defined on Y and such that $p_i(\beta) = 0$ for all $i \geq 3$. Hence, in fibred coordinates

$$(3.1) \quad \rho = L\omega_0 + \frac{\partial L}{\partial y_j^\sigma} \omega^\sigma \wedge \omega_j + g_{\sigma\nu}^{ij} \omega^\sigma \wedge \omega^\nu \wedge \omega_{ij},$$

where the functions $g_{\sigma\nu}^{ij}$ do not depend upon the y_l^κ ’s and satisfy the conditions

$$(3.2) \quad g_{\sigma\nu}^{ij} = -g_{\nu\sigma}^{ij}, \quad g_{\sigma\nu}^{ij} = -g_{\sigma\nu}^{ji}, \quad g_{\sigma\nu}^{ij} = g_{\nu\sigma}^{ji}.$$

Note that (3.2) mean that only

$$\binom{m}{2} \cdot \binom{n}{2} = \frac{1}{4}mn(m-1)(n-1)$$

of the $mn \times mn$ functions $g_{\sigma\nu}^{ij}$ are independent.

Theorem 1 [14]. *Let λ be a first-order Lagrangian, ρ its Lepagean equivalent as above. Assume that the matrix*

$$(3.3) \quad \left(\frac{\partial^2 L}{\partial y_i^\sigma \partial y_j^\nu} - 4g_{\sigma\nu}^{ij} \right)$$

with rows (respectively, columns) labelled by the pair (σ, i) (respectively, (ν, j)), is regular. Then ρ is regular. Moreover, every Hamilton extremal δ of ρ is of the form $\delta = J^1\gamma$, where γ is an extremal of λ .

The proof is obtained by a direct calculation from (2.9), and can be found in [14].

In view of the above theorem we have the following concept:

Definition 1. Let $W \subset J^1Y$ be an open set. A Lagrangian λ is called *regularizable on W* if it has a regular Lepagean equivalent ρ (3.1) defined on W . If $W = J^1Y$ we say that λ is *globally regularizable*. We say that λ is *locally regularizable* if it is regularizable in a neighbourhood of every point in J^1Y . The corresponding Lepagean equivalent ρ is then called a *(local) regularization* of λ .

Note that for regularizable Lagrangians, the problem of solving the Euler–Lagrange equations is *equivalent* to the problem of solving (appropriate) Hamilton equations.

An important class of regularizable Lagrangians is characterized by the following proposition.

Proposition 1. *Let $m \geq 2$. Then every Lagrangian L such that*

$$(3.4) \quad L = a + b_\sigma^j y_j^\sigma + c_{\sigma\nu}^{jk} y_j^\sigma y_k^\nu,$$

where a , b_σ^j and $c_{\sigma\nu}^{jk}$ are functions of (x^i, y^ρ) , is locally regularizable.

In particular, every first-order Lagrangian affine (respectively, quadratic) in the y_p^ρ 's is locally regularizable.

Proof. By assumption, $\partial^2 L / \partial y_i^\sigma \partial y_j^\nu$ are functions defined on an open subset of the total space Y . Let $\det(\partial^2 L / \partial y_i^\sigma \partial y_j^\nu) = 0$ at a point $x \in Y$. Since $m > 2$, one can find functions $g_{\sigma\nu}^{ij}$, antisymmetric in $(\sigma\nu)$ and (ij) , defined in a neighbourhood of x and such that at x the condition (3.3) is satisfied. However, since the determinant is a continuous function, the corresponding matrix must be nondegenerate in a neighbourhood U of x . With these g 's (independent of the y_p^ρ 's, as desired), the form $\rho = \theta_\lambda + g_{\sigma\nu}^{ij} \omega^\sigma \wedge \omega^\nu \wedge \omega_{ij}$ is a regularization of λ on $W = \pi_{1,0}^{-1}(U)$. $\square$

The following theorem provides a *generalization of Legendre transformation* to singular in the standard sense, but *regularizable Lagrangians*.

Theorem 2. *Consider a Lepagean form ρ given in a fibred chart (V, ψ) , $\psi = (x^i, y^\sigma)$ by (3.1) (3.2). Put*

$$(3.5) \quad p_\sigma^i = \frac{\partial L}{\partial y_i^\sigma} - 4g_{\sigma\nu}^{ij} y_j^\nu.$$

Let $x \in V_1 \subset J^1 Y$ be a point. If the matrix (3.3) is regular in a neighbourhood $W \subset V_1$ of x , then $(x^i, y^\sigma, y_j^\sigma) \rightarrow (x^i, y^\sigma, p_\sigma^j)$ is a coordinate transformation on W .

Proof. The above theorem follows immediately from the fact that

$$\frac{\partial p_\sigma^i}{\partial y_j^\nu} = \frac{\partial^2 L}{\partial y_i^\sigma \partial y_j^\nu} - 4g_{\sigma\nu}^{ij}.$$

□

Denote

$$(3.6) \quad H = -L + \frac{\partial L}{\partial y_i^\sigma} y_i^\sigma - 2g_{\sigma\nu}^{ij} y_i^\sigma y_j^\nu = -L + p_\sigma^i y_i^\sigma + 2g_{\sigma\nu}^{ij} y_i^\sigma y_j^\nu.$$

Now the Lepagean form (3.1), (3.2) reads

$$(3.7) \quad \begin{aligned} \rho &= L\omega_0 + \frac{\partial L}{\partial y_j^\sigma} \omega^\sigma \wedge \omega_j + g_{\sigma\nu}^{ij} \omega^\sigma \wedge \omega^\nu \wedge \omega_{ij} = \left(L - \frac{\partial L}{\partial y_i^\sigma} y_i^\sigma + 2g_{\sigma\nu}^{ij} y_i^\sigma y_j^\nu \right) \omega_0 \\ &+ \left(\frac{\partial L}{\partial y_i^\sigma} - 4g_{\sigma\nu}^{ij} y_j^\nu \right) dy^\sigma \wedge \omega_i + g_{\sigma\nu}^{ij} dy^\sigma \wedge dy^\nu \wedge \omega_{ij} \\ &= -H\omega_0 + p_\sigma^i dy^\sigma \wedge \omega_i + g_{\sigma\nu}^{ij} dy^\sigma \wedge dy^\nu \wedge \omega_{ij}. \end{aligned}$$

In analogy with the standard terminology we shall call H the *Hamiltonian* and p_σ^i *momenta* of the Lepagean form ρ (3.1), (3.2), and the corresponding coordinate transformation *Legendre transformation*; accordingly, the coordinates $(x^i, y^\sigma, p_\sigma^i)$ will be referred to as *Legendre coordinates* of ρ .

Corollary 1. *Let $(x^i, y^\sigma, p_\sigma^i)$ be the Legendre transformation associated with a Lepagean form ρ (3.1), (3.2). Then the matrix*

$$(3.8) \quad \left(\frac{\partial^2 H}{\partial p_\sigma^i \partial p_\nu^j} \right)$$

is regular and inverse to the matrix (3.3).

Proof. Explicit computations lead to

$$\begin{aligned} \frac{\partial L}{\partial y_j^\nu} &= p_\nu^j + 4g_{\nu\kappa}^{jk} y_k^\kappa, \quad \frac{\partial L}{\partial p_\sigma^i} = \frac{\partial L}{\partial y_j^\nu} \frac{\partial y_j^\nu}{\partial p_\sigma^i} = (p_\nu^j + 4g_{\nu\kappa}^{jk} y_k^\kappa) \frac{\partial y_j^\nu}{\partial p_\sigma^i}, \\ \frac{\partial H}{\partial p_\sigma^i} &= -\frac{\partial L}{\partial p_\sigma^i} + y_i^\sigma + p_\nu^j \frac{\partial y_j^\nu}{\partial p_\sigma^i} + 4g_{\kappa\nu}^{kj} y_k^\kappa \frac{\partial y_j^\nu}{\partial p_\sigma^i} = y_i^\sigma, \quad \frac{\partial^2 H}{\partial p_\sigma^i \partial p_\nu^j} = \frac{\partial y_i^\sigma}{\partial p_\nu^j}, \end{aligned}$$

proving the assertion. □

Expressing Hamilton p_2 -equations (2.9) in Legendre coordinates we get

$$(3.9) \quad \begin{aligned} \frac{\partial H}{\partial y^\sigma} &= -\frac{\partial p_\sigma^i}{\partial x^i} + 4\frac{\partial g_{\sigma\nu}^{ij}}{\partial x^j} \frac{\partial y^\nu}{\partial x^i} + 2\left(\frac{\partial g_{\kappa\nu}^{ij}}{\partial y^\sigma} + \frac{\partial g_{\sigma\kappa}^{ij}}{\partial y^\nu} + \frac{\partial g_{\nu\sigma}^{ij}}{\partial y^\kappa} \right) \frac{\partial y^\kappa}{\partial x^i} \frac{\partial y^\nu}{\partial x^j}, \\ \frac{\partial H}{\partial p_\sigma^i} &= \frac{\partial y^\sigma}{\partial x^i}, \end{aligned}$$

or, equivalently,

$$(3.10) \quad \begin{aligned} \frac{\partial H}{\partial y^\sigma} &= -\frac{\partial p_\sigma^i}{\partial x^i} + 4 \frac{\partial g_{\sigma\nu}^{ij}}{\partial x^j} \frac{\partial H}{\partial p_\nu^i} + 2 \left(\frac{\partial g_{\kappa\nu}^{ij}}{\partial y^\sigma} + \frac{\partial g_{\sigma\kappa}^{ij}}{\partial y^\nu} + \frac{\partial g_{\nu\sigma}^{ij}}{\partial y^\kappa} \right) \frac{\partial H}{\partial p_\kappa^i} \frac{\partial H}{\partial p_\nu^j}, \\ \frac{\partial H}{\partial p_\sigma^i} &= \frac{\partial y^\sigma}{\partial x^i}. \end{aligned}$$

Corollary 2. *If the n -form*

$$(3.11) \quad \eta = g_{\sigma\nu}^{ij} dy^\sigma \wedge dy^\nu \wedge \omega_{ij}$$

is closed then the above Hamilton p_2 -equations take the form

$$(3.12) \quad \frac{\partial H}{\partial y^\sigma} = -\frac{\partial p_\sigma^i}{\partial x^i}, \quad \frac{\partial H}{\partial p_\sigma^i} = \frac{\partial y^\sigma}{\partial x^i}.$$

Remark 1. Compared with Dedecker [3], we differ in both the definition of regularity and Legendre transformation. Dedecker’s regularity is weaker—Hamilton equations regular in his sense need not be equivalent with the Euler–Lagrange equations. Also, Legendre transformation is completely different: while Dedecker’s Legendre transformation is a map to a certain new space of higher dimension than that of the dynamical space, with unclear relations to regularity and to Lagrangian dynamics, Legendre transformation proposed above has similar properties as the Legendre transformation in classical mechanics (if the Lagrangian is regular in our sense, Legendre transformation becomes a local diffeomorphism of the space where the dynamics proceeds, providing a “canonical” form of the motion equations). Some further geometric properties of this Legendre transformation are clarified in [13].

4. SATELLITE LAGRANGIANS

We shall investigate the meaning of the 2-contact term in the Lepagean equivalent ρ (3.1) of a Lagrangian. Keeping notations introduced so far, we start with the following interesting assertions:

Lemma 1. *It holds (up to the projection $\pi_{1,0}$)*

$$(4.1) \quad \eta = \rho_{h(\eta)}.$$

Proof. Denote $h(\eta) = l \omega_0$; then obviously,

$$(4.2) \quad l = 2g_{\sigma\nu}^{ij} y_i^\sigma y_j^\nu,$$

and we obtain using (2.7) and (2.2)

$$(4.3) \quad \begin{aligned} \rho_{h(\eta)} &= l \omega_0 + \frac{\partial l}{\partial y_i^\sigma} \omega^\sigma \wedge \omega_i + \frac{1}{4} \frac{\partial^2 l}{\partial y_i^\sigma \partial y_j^\nu} \omega^\sigma \wedge \omega^\nu \wedge \omega_{ij} \\ &= 2g_{\sigma\nu}^{ij} y_i^\sigma y_j^\nu \omega_0 + 4g_{\sigma\nu}^{ij} y_j^\nu \omega^\sigma \wedge \omega_i + g_{\sigma\nu}^{ij} \omega^\sigma \wedge \omega^\nu \wedge \omega_{ij} \\ &= g_{\sigma\nu}^{ij} dy^\sigma \wedge dy^\nu \wedge \omega_{ij} = \eta. \end{aligned}$$

□

Lemma 2. *Put*

$$(4.4) \quad \bar{\lambda} = \lambda - h(\eta) = h(\rho - \eta), \quad i.e. \quad \bar{L} = L - l.$$

Then

$$(4.5) \quad \rho = \theta_{\bar{\lambda}} + \rho_{h(\eta)}, \quad i.e. \quad \theta_{\bar{\lambda}} = \rho - \eta.$$

Proof. Indeed, by (4.3) and (3.1), $\rho_{h(\eta)} = \theta_{h(\eta)} + p_2(\rho)$, hence, by (4.4),

$$\rho = \theta_{\lambda} + p_2(\rho) = \theta_{\lambda} - \theta_{h(\eta)} + \rho_{h(\eta)} = \theta_{\bar{\lambda}} + \rho_{h(\eta)}.$$

□

Note that by (4.5) and (3.7),

$$(4.6) \quad \theta_{\bar{\lambda}} = -H\omega_0 + p_{\sigma}^i dy^{\sigma} \wedge \omega_i.$$

Let us denote by $\tilde{p}_{\sigma}^j(L)$ and $\tilde{H}(L)$ the De Donder momenta and Hamiltonian associated with a Lagrangian L . Recall that

$$(4.7) \quad \tilde{p}_{\sigma}^i(L) = \frac{\partial L}{\partial y_i^{\sigma}}, \quad \tilde{H}(L) = -L + \tilde{p}_{\sigma}^i(L) y_i^{\sigma}$$

[4], [7]. Note that for our momenta and Hamiltonian (3.5) and (3.6) we obtain

$$(4.8) \quad p_{\sigma}^i = \tilde{p}_{\sigma}^i(L) - \tilde{p}_{\sigma}^j(l), \quad H = \tilde{H}(L) - \tilde{H}(l) = \tilde{H}(L) - l.$$

Consequently, we have the following lemma.

Lemma 3. *It holds*

$$(4.9) \quad H = \tilde{H}(\bar{L}), \quad p_{\sigma}^i = \tilde{p}_{\sigma}^i(\bar{L}).$$

Moreover, the regularity condition (3.3) is equivalent with the “standard” regularity condition (1.1) for $\bar{L}$, i.e., with

$$(4.10) \quad \det\left(\frac{\partial^2 \bar{L}}{\partial y_i^{\sigma} \partial y_k^{\nu}}\right) \neq 0.$$

In view of the above results we shall call the Lagrangian $h(\eta)$ a *satellite* of λ , and the Lagrangian $\bar{\lambda}$ a *dedonderization* of λ .

Now, from Lemma 1 and 2 we immediately obtain the following important result:

Proposition 2. *If the form η is closed then the Lagrangian $\bar{\lambda}$ is equivalent with λ , and $d\rho = d\theta_{\bar{\lambda}}$.*

In Section 3 we introduced *regularization* as a procedure to find for a Lagrangian appropriate *Hamilton p_2 -equations* (i.e., a certain “correction” to the Hamilton–De Donder equations of L) which are *equivalent* with the Euler–Lagrange equations, hence represent a suitable (unconstrained) alternative for solving the extremal problem. Now, taking into account all the above properties of satellite Lagrangians we conclude that regularization can be understood also *in a different way* as a procedure to find to a given Lagrangian an appropriate satellite in such a way that the Hamilton–De Donder equations of the new Lagrangian would be equivalent with the Euler–Lagrange equations:

Theorem 3. *Let λ be a regularizable Lagrangian. Then for every its (local) regularization ρ such that $d\eta = 0$, the Lagrangian $\bar{\lambda} = \lambda - h(\eta)$ is equivalent with λ , satisfies the “standard” regularity condition (4.10), and the Hamilton p_2 -equations of λ based upon ρ coincide with the Hamilton–De Donder equations of the Lagrangian $\bar{\lambda}$.*

5. EXAMPLES OF LEGENDRE TRANSFORMATIONS FOR FIRST-ORDER LAGRANGIANS

The above results can be directly applied to concrete Lagrangians. Let us consider two important cases: *Lagrangians affine in the first derivatives of the field variables* (in particular, the *Dirac field*), and the *electromagnetic field*.

5.1. Affine Lagrangians. Recall that by Proposition 1, if the fibre dimension m is at least 2, all Lagrangians affine in the first derivatives are locally regularizable and admit Legendre transformation introduced in Section 3. Assume

$$(5.1) \quad L = L_0 + L_\sigma^i y_i^\sigma$$

where L_0 and L_σ^i ($1 \leq i \leq n$, $1 \leq \sigma \leq m$) are functions of (x^j, y^ν) . Note that the De Donder momenta and Hamiltonian of (5.1) take the form $\tilde{p}_\sigma^j = L_\sigma^j$, $\tilde{H} = -L_0$, i.e., they are defined on an open subset of the *total space* Y , and the corresponding Hamilton equations must be treated as *constrained*, within the range of the Dirac theory of constrained systems (cf. eg. [6]). On the other hand, we get by (3.5) and (3.6),

$$(5.2) \quad p_\sigma^i = L_\sigma^i - 4g_{\sigma\nu}^{ij} y_j^\nu, \quad H = -L_0 - 2g_{\sigma\nu}^{ij} y_i^\sigma y_j^\nu,$$

where $(g_{\sigma\nu}^{ij})$ is a regular $(mn \times mn)$ -matrix. We can see that the domain of definition of the functions (5.2) is an open subset of $J^1 Y$; momenta are functions *affine* in the y_j^ν 's, and H (in Legendre coordinates) is a *polynomial of degree 2* in momenta. By Theorem 3, we should choose the $g_{\sigma\nu}^{ij}$ in such a way that the form η (3.11) be closed. Then the corresponding satellite Lagrangian is trivial, and Hamilton equations take the form (3.12).

Let us discuss in more detail the case $m = 2$, and $n = 2$ (respectively, $n = 4$).

(i) $n = \dim X = 2$. The conditions (3.2) on the $g_{\sigma\nu}^{ij}$'s mean that only one of these functions is independent, say, g_{12}^{12} . Denote $u = 4g_{12}^{12}$, and assume $u \neq 0$. The condition $d\eta = 0$, i.e., $du \wedge dy^1 \wedge dy^2 = 0$ means that $u = u(y^1, y^2)$.

As above, we consider Lepagean equivalents of the Lagrangian (5.1) in the form

$$(5.3) \quad \rho = (L_0 + L_\sigma^i y_i^\sigma) dx^1 \wedge dx^2 + L_\sigma^i \omega^\sigma \wedge \omega_i + u \omega^1 \wedge \omega^2$$

(where summation runs through $1 \leq i, \sigma \leq 2$). The regularity condition (3.3) reads

$$(5.4) \quad \det \begin{pmatrix} 0 & 0 & 0 & -u \\ 0 & 0 & u & 0 \\ 0 & u & 0 & 0 \\ -u & 0 & 0 & 0 \end{pmatrix} \neq 0,$$

and is clearly satisfied. Momenta become

$$(5.5) \quad p_1^1 = L_1^1 - uy_2^2, \quad p_2^1 = L_2^1 + uy_2^1, \quad p_1^2 = L_1^2 + uy_1^2, \quad p_2^2 = L_2^2 - uy_1^1.$$

Since the inverse to the Legendre transformation takes the form

$$(5.6) \quad y_1^1 = \frac{1}{u} (L_2^2 - p_2^2), \quad y_2^2 = \frac{1}{u} (L_1^1 - p_1^1), \quad y_1^2 = -\frac{1}{u} (L_1^2 - p_1^2), \quad y_2^1 = -\frac{1}{u} (L_2^1 - p_2^1),$$

the Hamiltonian in the Legendre coordinates reads

$$(5.7) \quad H = -L_0 + \frac{1}{u} \left(L_2^1 L_1^2 - L_1^1 L_2^2 + p_1^1 L_2^2 + p_2^2 L_1^1 - p_2^1 L_1^2 - p_1^2 L_2^1 - p_1^1 p_2^2 + p_2^1 p_1^2 \right).$$

Hamilton p_2 -equations in the Legendre coordinates take the form

$$(5.8) \quad \begin{aligned} \frac{\partial H}{\partial y^1} &= -\frac{\partial p_1^1}{\partial x^1} - \frac{\partial p_1^2}{\partial x^2}, & \frac{\partial H}{\partial y^2} &= -\frac{\partial p_2^1}{\partial x^1} - \frac{\partial p_2^2}{\partial x^2}, \\ \frac{\partial H}{\partial p_1^1} &= \frac{\partial y^1}{\partial x^1}, & \frac{\partial H}{\partial p_2^1} &= \frac{\partial y^1}{\partial x^2}, & \frac{\partial H}{\partial p_1^2} &= \frac{\partial y^2}{\partial x^1}, & \frac{\partial H}{\partial p_2^2} &= \frac{\partial y^2}{\partial x^2}. \end{aligned}$$

Note that for every fixed $u \neq 0$ we have obtained Hamilton equations *equivalent* with the Euler-Lagrange equations.

As an illustration, let us consider the *Dirac field*. In this case we have $X = R^2$, $Y = R^2 \times R^2$, i.e. $J^1 Y = R^2 \times R^2 \times R^4$, with the global coordinates denoted by $(x^\mu, \psi, \bar{\psi}, \partial_\mu \psi, \partial_\mu \bar{\psi})$, $\mu = 1, 2$, Lagrangian L takes the form

$$(5.9) \quad L = \frac{i}{2} (\bar{\psi} \gamma^\mu \partial_\mu \psi + \partial_\mu \bar{\psi} \gamma^\mu \psi) - \bar{\psi} m \psi,$$

and it is apparently degenerate in the sense of the standard regularity condition (1.1). However, for every nonzero function $u(\psi, \bar{\psi})$, the form ρ (5.3) is a *global regularization* of L . Computing the corresponding *satellite Lagrangian* for (5.9) we get

$$(5.10) \quad l = -u \epsilon^{\mu\nu} \partial_\mu \bar{\psi} \partial_\nu \psi,$$

where $\epsilon^{\mu\nu}$ is the Levi-Civita symbol. Now the Lagrangian

$$(5.11) \quad \bar{L} = \frac{i}{2} (\bar{\psi} \gamma^\mu \partial_\mu \psi + \partial_\mu \bar{\psi} \gamma^\mu \psi) - \bar{\psi} m \psi + u \epsilon^{\mu\nu} \partial_\mu \bar{\psi} \partial_\nu \psi$$

is a dedonderization of the Dirac field Lagrangian, which is *regular* in the standard sense. Accordingly, Hamilton–De Donder equations of (5.11) are equivalent with the Dirac field equations, and in this sense, the (standard) De Donder Hamiltonian and momenta of (5.11) (which, however, are precisely the functions H and p ’s given by (5.7) and (5.5)), should represent possibly a better physical alternative for (unconstrained) quantization of the Dirac field. Explicitly, the “new” Hamiltonian reads by (4.8)

$$(5.12) \quad H = u \epsilon^{\mu\nu} \partial_\mu \bar{\psi} \partial_\nu \psi + \bar{\psi} m \psi.$$

Note that in the above formulas the most simple admissible choice is $u \neq 0$ a *constant* function.

(ii) $n = \dim X = 4$. In this case we get 6 independent functions $g_{\sigma\nu}^{ij}$. Denote

$$(5.13) \quad u_1 = 4g_{12}^{12}, \quad u_2 = 4g_{12}^{13}, \quad u_3 = 4g_{12}^{14}, \quad u_4 = 4g_{12}^{23}, \quad u_5 = 4g_{12}^{24}, \quad u_6 = 4g_{12}^{34}.$$

The matrix (3.3) takes the form

$$(5.14) \quad \begin{pmatrix} 0 & -M \\ M & 0 \end{pmatrix}, \quad \text{where} \quad M = \begin{pmatrix} 0 & u_1 & u_2 & u_3 \\ -u_1 & 0 & u_4 & u_5 \\ -u_2 & -u_4 & 0 & u_6 \\ -u_3 & -u_5 & -u_6 & 0 \end{pmatrix}.$$

We can see that for any choice of functions $u_k(x^i, y^\sigma)$, $1 \leq k \leq 6$, such that $\det M \neq 0$ we obtain a regular Hamilton p_2 -theory, based upon the Lepagean form

$$(5.15) \quad \begin{aligned} \rho = & (L_0 + L_\sigma^i y_i^\sigma) dx^1 \wedge dx^2 \wedge dx^3 \wedge dx^4 + L_\sigma^i \omega^\sigma \wedge \omega_i \\ & + u_1 \omega^1 \wedge \omega^2 \wedge \omega_{12} + u_2 \omega^1 \wedge \omega^2 \wedge \omega_{13} + u_3 \omega^1 \wedge \omega^2 \wedge \omega_{14} \\ & + u_4 \omega^1 \wedge \omega^2 \wedge \omega_{23} + u_5 \omega^1 \wedge \omega^2 \wedge \omega_{24} + u_6 \omega^1 \wedge \omega^2 \wedge \omega_{34}. \end{aligned}$$

Expressing the corresponding *satellite Lagrangian* for the Dirac field explicitly, we easily obtain

$$(5.16) \quad l = - \sum_{\mu, \nu} u_{(\mu, \nu)} \epsilon^{\mu\nu} \partial_\mu \bar{\psi} \partial_\nu \psi$$

where $u_{(\mu, \nu)} = u_{(\nu, \mu)}$ and the notation $u_1 = u_{(1,2)}$, $u_2 = u_{(1,3)}$, $u_3 = u_{(1,4)}$, $u_4 = u_{(2,3)}$, $u_5 = u_{(2,4)}$, $u_6 = u_{(3,4)}$ is used. “Corrected” momenta can now be obtained by a short routine calculation from (3.5), and the Hamiltonian takes by (4.8) the form $H = l + \bar{\psi} m \psi$. In comparison with the usual formulas they differ by additional terms—(De Donder) momenta and Hamiltonian of the satellite (5.16) of the Lagrangian (5.9).

Note that on the fibred manifold $R^4 \times R^2 \rightarrow R^4$ the most simple *global* regularization of the Dirac Lagrangian is obtained for u_i , $1 \leq i \leq 6$, constant functions; by (5.13) some of them may even equal zero (a simple choice is e.g. $u_3, u_4 \neq 0$, $u_1 = u_2 = u_5 = u_6 = 0$).

5.2. The electromagnetic field. For the electromagnetic field Lagrangian

$$(5.17) \quad L = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} = \frac{1}{2} (y_\nu^\sigma y_\sigma^\nu - \mathfrak{g}^{\sigma\nu} \mathfrak{g}_{\mu\rho} y_\sigma^\mu y_\nu^\rho)$$

(where $F_{\mu\nu} = A_{\mu, \nu} - A_{\nu, \mu}$, $(\mathfrak{g}_{\sigma\nu})$ denotes the Lorentz metric, $\mathfrak{g}_{\sigma\nu} = 0$ for $\sigma \neq \nu$, $-\mathfrak{g}_{11} = \mathfrak{g}_{22} = \mathfrak{g}_{33} = \mathfrak{g}_{44} = 1$, and $y^\sigma = \mathfrak{g}^{\sigma\nu} A_\nu$), the standard regularity condition (1.1) gives that L is degenerate. For example, in the $n = 2$ case, the De Donder momenta and Hamiltonian read

$$(5.18) \quad \tilde{p}_1^1 = \tilde{p}_2^2 = 0, \quad \tilde{p}_2^1 = \tilde{p}_1^2 = y_2^1 + y_1^2, \quad \tilde{H} = -\frac{1}{2} (y_2^1 + y_1^2)^2 + \tilde{p}_2^1 y_1^2 + \tilde{p}_1^2 y_2^1.$$

Hence, momenta are not independent, and the corresponding Hamilton equations must be treated as *constrained*. However, as we proved in Sec. 3, Lagrangian (5.17) is regularizable (and admits many global regularizations). Let us choose one of them and compute the corresponding Hamiltonian and momenta. Put

$$(5.19) \quad g_{\sigma\nu}^{\alpha\beta} = \frac{\partial^2 L}{\partial y_\alpha^\sigma \partial y_\beta^\nu} - \frac{\partial^2 L}{\partial y_\beta^\sigma \partial y_\alpha^\nu}$$

(apparently, these g ’s do not depend upon the y_i^σ ’s, and satisfy (3.2), as desired). The Lepagean equivalent (3.1) now reads,

$$(5.20) \quad \rho = L \omega_0 + \frac{\partial L}{\partial y_\alpha^\sigma} \omega^\sigma \wedge \omega_\alpha + \left(\frac{\partial^2 L}{\partial y_\alpha^\sigma \partial y_\beta^\nu} - \frac{\partial^2 L}{\partial y_\beta^\sigma \partial y_\alpha^\nu} \right) \omega^\sigma \wedge \omega^\nu \wedge \omega_{\alpha\beta},$$

and the regularity condition (3.3) leads to checking regularity of the following matrix

$$(5.21) \quad \left(4 \frac{\partial^2 L}{\partial y_\beta^\sigma \partial y_\alpha^\nu} - 3 \frac{\partial^2 L}{\partial y_\alpha^\sigma \partial y_\beta^\nu} \right).$$

(i) Let $X = R^2$. Then we have $Y = R^2 \times R^2$, i.e. $m = 2$, and the Lagrangian (5.17) reads

$$L = \frac{1}{2} (y_2^1 + y_1^2)^2.$$

For the matrix (5.21) we obtain

$$\begin{pmatrix} 0 & 0 & 0 & 4 \\ 0 & 1 & -3 & 0 \\ 0 & -3 & 1 & 0 \\ 4 & 0 & 0 & 0 \end{pmatrix},$$

i.e., it is *regular*. Consequently, the related Hamilton p_2 -equations are *equivalent* with the Maxwell equations. For the momenta we easily obtain

$$(5.22) \quad p_1^1 = 4y_2^2, \quad p_2^1 = -3y_2^1 + y_1^2, \quad p_2^2 = 4y_1^1, \quad p_1^2 = y_2^1 - 3y_1^2.$$

The inverse transformation to the Legendre transformation exists and takes the form

$$y_1^1 = \frac{1}{4}p_2^2, \quad y_2^1 = -\frac{3}{8}p_2^1 - \frac{1}{8}p_1^2, \quad y_2^2 = \frac{1}{4}p_1^1, \quad y_1^2 = -\frac{1}{8}p_2^1 - \frac{3}{8}p_1^2.$$

The Hamiltonian in the Legendre coordinates reads

$$(5.23) \quad H = \frac{1}{4}p_1^1p_2^2 - \frac{3}{8}p_2^1p_1^2 - \frac{1}{16}(p_2^1)^2 - \frac{1}{16}(p_1^2)^2.$$

(ii) Let $X = R^4$. We have $m = 4$, and the Lagrangian (5.17) takes the form

$$(5.24) \quad \begin{aligned} L = & \frac{1}{2} (y_2^1 + y_1^2)^2 + \frac{1}{2} (y_3^1 + y_1^3)^2 + \frac{1}{2} (y_4^1 + y_1^4)^2 \\ & - \frac{1}{2} (y_3^2 - y_2^3)^2 - \frac{1}{2} (y_4^2 - y_2^4)^2 - \frac{1}{2} (y_4^3 - y_3^4)^2. \end{aligned}$$

The matrix (5.21) becomes

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 \\ 0 & 1 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 \\ 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 4 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -3 & 0 & 0 & 0 & -1 & 0 \\ 4 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 4 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and one can easily check that it is regular. For the momenta we get

$$(5.25) \quad \begin{aligned} p_1^1 &= 4y_2^2 + 4y_3^3 + 4y_4^4, & p_1^2 &= y_2^1 - 3y_1^2, & p_1^3 &= y_3^1 - 3y_1^3, & p_1^4 &= y_4^1 - 3y_1^4, \\ p_2^2 &= 4y_1^1 + 4y_3^3 + 4y_4^4, & p_2^1 &= y_1^2 - 3y_2^1, & p_2^3 &= -y_3^2 - 3y_2^3, & p_2^4 &= -y_4^2 - 3y_2^4, \\ p_3^3 &= 4y_1^1 + 4y_2^2 + 4y_4^4, & p_3^1 &= y_1^3 - 3y_3^1, & p_3^2 &= -y_2^3 - 3y_3^2, & p_3^4 &= -y_4^3 - 3y_3^4, \\ p_4^4 &= 4y_1^1 + 4y_2^2 + 4y_3^3, & p_4^1 &= y_1^4 - 3y_4^1, & p_4^2 &= -y_2^4 - 3y_4^2, & p_4^3 &= -y_3^4 - 3y_4^3, \end{aligned}$$

and the Hamiltonian in the Legendre coordinates takes the form

$$(5.26) \quad \begin{aligned} H &= -\frac{1}{12} \left((p_1^1)^2 + (p_2^2)^2 + (p_3^3)^2 + (p_4^4)^2 - p_1^1 p_2^2 - p_1^1 p_3^3 - p_1^1 p_4^4 - p_2^2 p_3^3 - p_2^2 p_4^4 - p_3^3 p_4^4 \right) \\ &\quad - \frac{1}{16} \left((p_2^1)^2 + (p_1^2)^2 + (p_3^1)^2 + (p_1^3)^2 + (p_4^1)^2 + (p_1^4)^2 \right. \\ &\quad \left. - (p_3^2)^2 - (p_2^3)^2 - (p_4^2)^2 - (p_2^4)^2 - (p_4^3)^2 - (p_3^4)^2 \right) \\ &\quad - \frac{3}{8} \left(p_2^1 p_1^2 + p_3^1 p_1^3 + p_4^1 p_1^4 + p_3^2 p_2^3 + p_4^2 p_2^4 + p_4^3 p_3^4 \right). \end{aligned}$$

Let us compute the corresponding *satellite Lagrangian* for (5.17). Since

$$(5.27) \quad g_{\sigma\nu}^{\alpha\beta} = \delta_\nu^\alpha \delta_\sigma^\beta - \delta_\nu^\beta \delta_\sigma^\alpha,$$

we obtain

$$(5.28) \quad l = 2(A_\nu^\mu A_\mu^\nu - A_\mu^\mu A_\nu^\nu) = 2(\text{Tr}(A'^2) - (\text{Tr } A')^2),$$

where A' denotes the matrix $(\partial_\alpha A^\beta)$. Now, the dedonderization Lagrangian and the “new” Hamiltonian for the electromagnetic field read

$$(5.29) \quad \bar{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + 2(A_\mu^\mu A_\nu^\nu - A_\nu^\mu A_\mu^\nu), \quad H = \tilde{H} + 2(A_\mu^\mu A_\nu^\nu - A_\nu^\mu A_\mu^\nu).$$

Remark 2. Note that the Lepagean equivalent ρ_λ (2.7) is *not* an appropriate regularization of the electromagnetic field. Indeed, in this case the matrix (3.3) takes the form

$$\left(\frac{\partial^2 L}{\partial y_\beta^\sigma \partial y_\alpha^\nu} + \frac{\partial^2 L}{\partial y_\alpha^\sigma \partial y_\beta^\nu} \right),$$

and it is singular for the Lagrangian (5.17).

REFERENCES

1. D. E. Betounes, *Extension of the classical Cartan form*, Phys. Rev. D 29 (1984), 599–606.
2. F. Cantrijn, L.A. Ibort and M. de León, *Hamiltonian structures on multisymplectic manifolds*, Rend. Sem. Mat. Univ. Pol. Torino 54 (1996) 225–236.
3. P. Dedecker, *On the generalization of symplectic geometry to multiple integrals in the calculus of variations*, in: Lecture Notes in Math. 570 (Springer, Berlin, 1977) 395–456.
4. Th. De Donder, *Théorie Invariantive du Calcul des Variations*, Gauthier–Villars, Paris, 1930.
5. G. Giachetta, L. Mangiarotti and G. Sardanashvily, *New Lagrangian and Hamiltonian Methods in Field Theory*, World Scientific, Singapore, 1997.

6. G. Giachetta, L. Mangiarotti and G. Sardanashvily, *Covariant Hamilton equations for field theory*, J. Phys. A: Math. Gen. 32 (1999) 6629–6642.
7. H. Goldschmidt and S. Sternberg, *The Hamilton–Cartan formalism in the calculus of variations*, Ann. Inst. Fourier, Grenoble 23 (1973), 203–267.
8. M. J. Gotay, *A multisymplectic framework for classical field theory and the calculus of variations, I. Covariant Hamiltonian formalism*, in: Mechanics, Analysis and Geometry: 200 Years After Lagrange, M. Francaviglia and D. D. Holm, eds. (North Holland, Amsterdam, 1990) 203–235.
9. D. Krupka, *A geometric theory of ordinary first order variational problems in fibred manifolds. I. Critical sections, II. Invariance*, J. Math. Anal. Appl. 49 (1975), 180–206; 469–476.
10. D. Krupka, *A map associated to the Lepagean forms of the calculus of variations in fibred manifolds*, Czechoslovak Math. J. 27 (1977), 114–118.
11. D. Krupka, *On the higher order Hamilton theory in fibred spaces*, in: Geometrical Methods in Physics, Proc. Conf. Diff. Geom. Appl., Nové Město na Moravě, 1983, D. Krupka, ed. (J.E. Purkyně University, Brno, Czechoslovakia, 1984) 167–183.
12. D. Krupka and O. Štěpánková, *On the Hamilton form in second order calculus of variations*, in: Geometry and Physics, Proc. Int. Meeting, Florence, Italy, 1982, M. Modugno, ed. (Pitagora Ed., Bologna, 1983) 85–101.
13. O. Krupková, *Hamiltonian field theory revisited: A geometric approach to regularity*, Preprint GA 11/2000 (Silesian University, Opava, 2000) 19pp; submitted to Proc. Colloq. Diff. Geom., Debrecen, July 2000.
14. O. Krupková and D. Smetanová, *On regularization of variational problems in first-order field theory*, Rend. Circ. Mat. Palermo Suppl., to appear.
15. M. de León and P.R. Rodrigues, *Generalized Classical Mechanics and Field Theory*, North-Holland, Amsterdam, 1985.
16. J.E. Marsden and S. Shkoller, *Multisymplectic geometry, covariant Hamiltonians, and water waves*, Math. Proc. Camb. Phil. Soc. 125 (1999) 553–575.
17. D.J. Saunders, *The regularity of variational problems*, Contemporary Math., 132 (1992) 573–593.
18. D. Smetanová, *On Hamilton p_2 -equations in second order field theory*, Proc. Colloq. Diff. Geom., Debrecen, Hungary, July 2000, submitted.